

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 2

✓
Please amend the claims as follows:

1. (Amended) A method of optimizing therapeutic efficacy [of 6-mercaptopurine drug] for treatment of an immune-mediated gastrointestinal disorder, comprising:

(a) administering a [6-mercaptopurine] drug to a subject having said immune-mediated gastrointestinal disorder, wherein said drug provides 6-thioguanine to said subject; and

(b) determining a level of 6-thioguanine in said subject having said immune-mediated gastrointestinal disorder,

wherein a level of 6-thioguanine less than a level corresponding to about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of [6-mercaptopurine] said drug subsequently administered to said subject and

wherein a level of 6-thioguanine greater than a level corresponding to about 400 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of [6-mercaptopurine] said drug subsequently administered to said subject.

7. (Amended) A method of reducing toxicity associated with [6-mercaptopurine drug] treatment of an immune-mediated gastrointestinal disorder, comprising:

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 3

(a) administering a [6-mercaptopurine] drug to a subject having said immune-mediated gastrointestinal disorder, wherein said drug provides 6-thioguanine to said subject; [and]

(b) determining a level of [a 6-mercaptopurine metabolite] 6-thioguanine in said subject having said immune-mediated gastrointestinal disorder[,]; and

(c) determining a level of 6-methyl-mercaptopurine in said subject having said immune-mediated gastrointestinal disorder,

wherein a level of [said 6-mercaptopurine metabolite] 6-thioguanine greater than [a predetermined toxic level of said 6-mercaptopurine metabolite] about 400 pmol per 8×10^8 red blood cells or a level of 6-methyl-mercaptopurine greater than about 7000 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of [6-mercaptopurine] said drug subsequently administered to said subject, thereby reducing toxicity associated with [6-mercaptopurine] said drug treatment of said immune-mediated gastrointestinal disorder.

B3 11/13. (Amended) The method of claim [11] 7, wherein said toxicity associated with [6-mercaptopurine] said drug treatment is hematologic toxicity.

B4 12/16. (Amended) The method of claim [14] 7, wherein said toxicity associated with [6-mercaptopurine] said drug treatment is hepatic toxicity.

B1

B

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 4

B4
concl'd

13

17. (Amended) The method of claim 7, wherein said level of [6-mercaptopurine metabolite] 6-thioguanine and said level of 6-methyl-mercaptopurine each is determined in red blood cells.

19. (Amended) A method of optimizing therapeutic efficacy and reducing toxicity associated with [6-mercaptopurine drug] treatment of an immune-mediated gastrointestinal disorder, comprising:

Sub 3

(a) administering a [6-mercaptopurine] drug to a subject having said immune-mediated gastrointestinal disorder, wherein said drug provides 6-thioguanine to said subject;

(b) determining a level of 6-thioguanine in said subject having said immune-mediated gastrointestinal disorder; and

(c) determining a level of 6-methyl-mercaptopurine in said subject having said immune-mediated gastrointestinal disorder,

wherein a level of 6-thioguanine less than [a predetermined minimal therapeutic level] about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of [6-mercaptopurine] said drug subsequently administered to said subject, thereby increasing therapeutic efficacy,

B

B

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 5

Sub 3
7/5
comp'd

wherein a level of 6-thioguanine greater than [a predetermined toxic level of 6-thioguanine] about 400 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of [6-mercaptopurine] said drug subsequently administered to said subject, thereby reducing toxicity associated with [6-mercaptopurine] said drug treatment of said immune-mediated gastrointestinal disorder, and

wherein a level of 6-methyl-mercaptopurine greater than [a predetermined toxic level of 6-methyl-mercaptopurine] about 7000 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of [6-mercaptopurine] said drug subsequently administered to said subject, thereby reducing toxicity associated with [6-mercaptopurine] said drug treatment of said immune-mediated gastrointestinal disorder.

In claim 29, line 2, please delete "6-mercaptopurine" and insert therefor --said--.

Sub 4

30. (Amended) A method of optimizing therapeutic efficacy of [6-mercaptopurine drug] treatment of a non-IBD autoimmune disease, comprising:

(a) administering a [6-mercaptopurine] drug to a subject having said non-IBD autoimmune disease, wherein said drug provides 6-thioguanine to said subject; and

(b) determining a level of 6-thioguanine in said subject having said non-IBD autoimmune disease,

B

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 6

Sub C4
B
could

wherein a level of 6-thioguanine less than [a minimal therapeutic level] about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of [6-mercaptopurine] said drug subsequently administered to said subject and

wherein a level of 6-thioguanine greater than [a level corresponding to a predetermined toxic level] about 400 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of [6-mercaptopurine] said drug subsequently administered to said subject.

In claim 33, line 2, please delete "6-mercaptopurine metabolite" and insert therefor --6-thioguanine--.

Sub C5
B

35. (Amended) A method of optimizing therapeutic efficacy and reducing toxicity associated with [6-mercaptopurine drug] treatment of an immune-mediated gastrointestinal disorder, comprising:

(a) administering a [6-mercaptopurine] drug to a subject having said immune-mediated gastrointestinal disorder, wherein said drug provides 6-thioguanine to said subject; [and]

(b) determining a level of [a 6-mercaptopurine metabolite] 6-thioguanine in said subject having said immune-mediated gastrointestinal disorder[,]; and

B

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 7

(c) determining a level of 6-methyl-mercaptopurine in said subject having said immune-mediated gastrointestinal disorder,

wherein a level of [said 6-mercaptopurine metabolite] 6-thioguanine less than [a predetermined minimal therapeutic level] about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of [6-mercaptopurine] said drug subsequently administered to said subject, thereby increasing therapeutic efficacy, and

wherein a level of [said 6-mercaptopurine metabolite] 6-thioguanine greater than [a predetermined toxic level of said 6-mercaptopurine metabolite] about 400 pmol per 8×10^8 red blood cells or a level of 6-methyl-mercaptopurine greater than about 7000 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of [6-mercaptopurine] said drug subsequently administered to said subject, thereby reducing toxicity associated with [6-mercaptopurine] said drug treatment of said immune-mediated gastrointestinal disorder.

29 44. (Amended) The method of claim 25, wherein said level of [said 6-mercaptopurine metabolite] 6-thioguanine and said level of 6-methyl-mercaptopurine each is determined in red blood cells.

In claim 46, line 2, please delete "6-mercaptopurine" and insert therefor --said--.

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 8

✓
Please add the following new claims.

~~32~~
~~47~~. (New) The method of claim 1, wherein said drug is selected from the group consisting of 6-mercaptopurine, azathioprine, 6-thioguanine, and 6-methylmercaptopurine riboside.

~~33~~
~~48~~. (New) The method of claim 7, wherein said drug is selected from the group consisting of 6-mercaptopurine, azathioprine, 6-thioguanine, and 6-methylmercaptopurine riboside.

~~34~~
~~49~~. (New) The method of claim ~~15~~
~~19~~, wherein said drug is selected from the group consisting of 6-mercaptopurine, azathioprine, 6-thioguanine, and 6-methylmercaptopurine riboside.

~~35~~
~~50~~. (New) The method of claim ~~22~~
~~30~~, wherein said drug is selected from the group consisting of 6-mercaptopurine, azathioprine, 6-thioguanine, and 6-methylmercaptopurine riboside.

~~36~~
~~51~~. (New) The method of claim ~~25~~
~~35~~, wherein said drug is selected from the group consisting of 6-mercaptopurine, azathioprine, 6-thioguanine, and 6-methylmercaptopurine riboside.

52. (New) A method of optimizing therapeutic efficacy of treatment of a non-IBD autoimmune disease, comprising:

(a) administering a drug to a subject having said non-IBD autoimmune disease, wherein said drug provides 6-thioguanine to said subject;

Inventors: Seidman and Théorêt
Serial No.: 09/288,344
Filed: April 8, 1999
Page 9

(b) determining a level of 6-thioguanine in said subject having said non-IBD autoimmune disease; and

(c) determining a level of 6-methyl-mercaptopurine in said subject having said non-IBD autoimmune disease,

wherein a level of 6-thioguanine less than about 230 pmol per 8×10^8 red blood cells indicates a need to increase the amount of said drug subsequently administered to said subject and

wherein a level of 6-thioguanine greater than about 400 pmol per 8×10^8 red blood cells or a level of 6-methyl-mercaptopurine greater than about 7000 pmol per 8×10^8 red blood cells indicates a need to decrease the amount of said drug subsequently administered to said subject, thereby reducing toxicity associated with said drug treatment of said non-IBD autoimmune disease.

53. (New) The method of claim 52, wherein said level of 6-thioguanine and said level of 6-methyl-mercaptopurine each is determined in red blood cells.

54. (New) The method of claim 53, wherein said level is determined using high pressure liquid chromatography.--

B